

SODIUM CHLORIDE- sodium chloride injection, solution
B. Braun Medical Inc.

0.9% Sodium Chloride Injection USP

Partial Fill

DESCRIPTION

Each mL of 0.9% Sodium Chloride Injection USP contains:

Sodium Chloride USP 9 mg; Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5–7.0)

Calculated Osmolarity: 310 mOsmol/liter

Concentration of Electrolytes (mEq/100 mL): Sodium 15.4 Chloride 15.4

This solution is sterile, nonpyrogenic, isotonic and contains no bacteriostatic or antimicrobial agents.

The formula of the active ingredient is:

Ingredient	Molecular Formula	Molecular Weight
Sodium Chloride USP	NaCl	58.44

Not made with natural rubber latex, PVC or DEHP.

The plastic container is a copolymer of ethylene and propylene formulated and developed for parenteral drugs. The copolymer contains no plasticizers and exhibits virtually no leachability. The safety of the plastic container has been confirmed by biological evaluation procedures. The material passes Class VI testing as specified in the U.S. Pharmacopeia for Biological Tests — Plastic Containers. These tests have shown that the container is nontoxic and biologically inert.

The container/solution unit is a closed system and is not dependent upon entry of external air during administration. The container has two ports, one is for the intravenous administration set and the other is a medication addition site. Each is covered by a tamperproof barrier. Refer to the **Directions for Use** of the container to properly identify the ports.

No vapor barrier is necessary.

CLINICAL PHARMACOLOGY

0.9% Sodium Chloride Injection USP provides electrolytes and is a source of water for hydration. It is capable of inducing diuresis depending on the clinical condition of the patient.

Sodium, the major cation of the extracellular fluid, functions primarily in the control of water distribution, fluid balance, and osmotic pressure of body fluids. Sodium is also associated with chloride and bicarbonate in the regulation of the acid-base equilibrium of body fluid.

Chloride, the major extracellular anion, closely follows the metabolism of sodium, and changes in the acid-base balance of the body are reflected by changes in the chloride concentration.

INDICATIONS AND USAGE

This intravenous solution is indicated for use in adults and pediatric patients as a source of electrolytes and water for hydration.

This product is designed for use as a diluent and delivery system for intermittent intravenous administration of compatible drug additives. Consult prescribing information for **INDICATIONS AND USAGE** of drug additives to be administered in this manner.

CONTRAINDICATIONS

This solution is contraindicated where the administration of sodium or chloride could be clinically detrimental.

WARNINGS

Solutions containing sodium ions should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency, and in clinical states in which there is sodium retention with edema. In patients with diminished renal function, administration of solutions containing sodium ions may result in sodium retention.

PRECAUTIONS

General

Clinical evaluation and periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation. Significant deviations from normal concentrations may require tailoring of the electrolyte pattern, in this or an alternative solution(s).

Sodium-containing solutions should be administered with caution to patients receiving corticosteroids or corticotropin, or to other salt-retaining patients. Care should be exercised in administering solutions containing sodium to patients with renal or cardiovascular insufficiency, with or without congestive heart failure, particularly if they are postoperative or elderly. See **PRECAUTIONS, *Geriatric Use***.

To minimize the risk of possible incompatibilities arising from mixing this solution with other additives that may be prescribed, the final infusate should be inspected for cloudiness or precipitation immediately after mixing, prior to administration, and periodically during administration.

Laboratory Tests

Periodic laboratory determinations are necessary to monitor changes in fluid balance, electrolyte concentrations, and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient warrants such evaluation.

Drug Interactions

Sodium-containing solutions should be administered with caution to patients receiving corticosteroids or corticotrophin.

Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Studies have not been conducted to evaluate additional drug/drug or drug/food interactions with Sodium Chloride Injection USP.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies with 0.9% Sodium Chloride Injection USP have not been performed to evaluate carcinogenic potential, mutagenic potential or effects on fertility.

Pregnancy

Teratogenic Effects

Pregnancy Category C

Animal reproduction studies have not been conducted with 0.9% Sodium Chloride Injection USP. It is also not known whether 0.9% Sodium Chloride Injection USP can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. 0.9% Sodium Chloride Injection USP should be given to a pregnant woman only if clearly needed.

Labor and Delivery

Studies have not been conducted to evaluate the effects of Sodium Chloride Injection USP on labor and delivery. As reported in the literature, sodium chloride solutions have been administered during labor and delivery. Caution should be exercised, and the fluid balance, glucose and electrolyte concentrations and acid-base balance, of both mother and fetus should be evaluated periodically or whenever warranted by the condition of the patient or fetus.

Nursing Mothers

Because many drugs are excreted in human milk, caution should be exercised when 0.9% Sodium Chloride Injection USP is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of sodium chloride injections in pediatric patients have not been established by adequate and well controlled trials, however, the use of Sodium Chloride solutions in the pediatric population is referenced in the medical literature. All Warnings, Precautions and Adverse Reactions described in this label apply to pediatric patients.

Geriatric Use

An evaluation of current literature revealed no clinical experience identifying differences in response between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

Reactions which may occur because of the solution or the technique of administration include febrile response, infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection, extravasation, and hypervolemia.

The physician should also be alert to the possibility of adverse reactions to drug additives diluted and administered from the plastic partial fill container. Prescribing information for drug additives to be administered in this manner should be consulted.

Symptoms may result from an excess or deficit of one or more of the ions present in the solution; therefore, frequent monitoring of electrolyte levels is essential.

Hypernatremia may be associated with edema and exacerbation of congestive heart failure due to the retention of water, resulting in an expanded extracellular fluid volume.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures, and save the remainder of the fluid for examination if deemed necessary.

OVERDOSAGE

In the event of a fluid or solute overload during parenteral therapy, reevaluate the patient's condition, and institute appropriate corrective treatment.

DOSAGE AND ADMINISTRATION

This solution is for intravenous use only.

Do not use plastic container in series connection.

If administration is controlled by a pumping device, care must be taken to discontinue pumping action before the container runs dry or air embolism may result.

This solution is intended for intravenous administration using sterile equipment. It is recommended that intravenous administration apparatus be replaced at least once every 24 hours.

Use only if solution is clear and container and seals are intact.

As directed by a physician. Dosage is dependent upon the age, weight, and clinical condition of the patient as well as laboratory determinations.

There is no specific pediatric dose. The dose is dependent on weight, clinical condition, and laboratory results. Follow recommendations of appropriate pediatric reference text. See **PRECAUTIONS, Pediatric Use.**

When using this product as a diluent or vehicle for administration of drug additives, consult the prescribing information of the drug to be used.

Addition of medication should be accomplished using aseptic technique in order to assure sterility.

Physicochemical studies have shown that the container and solution can withstand freezing.

Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

0.9% Sodium Chloride Injection USP is supplied sterile and nonpyrogenic in partial fill polyolefin containers. The 100/150 mL product is packaged 64 per case. The 50/100 mL product is packaged 84 per case. The 25/100 mL product is packaged 116 per case.

NDC	REF	Fill/Container (mL)
0.9% Sodium Chloride Injection USP		
0264-1800-36	S8004-5410	25/100
0264-1800-31	S8004-5384	50/100
0264-1800-32	S8004-5264	100/150

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended that the product be stored at room temperature (25°C); however, brief exposure up to 40°C does not adversely affect the product.

Rx only

Revised: October 2016

PAB is a registered trademark of B. Braun Medical Inc.

Directions For Use of PAB® Container

Partial Additive Bag

Aseptic technique is required.

Caution: Before use, perform the following checks: Read the label. Ensure solution is the one ordered and is within the expiration date.

Inspect the solution in good light for cloudiness, haze or particulate matter; check the container for leakage or damage. Any container which is suspect should not be used.

Use only if solution is clear and container and seals are intact. Single dose container. Discard unused portion. Consult Package Insert for complete product information.


The physician should also be alert to the possibility of adverse reactions to drug additives diluted and administered from the plastic partial fill container. Prescribing information for drug additives to be administered in this manner should be consulted.

Do not use plastic container in series connection.

This solution is intended for intravenous administration using sterile equipment. It is recommended that intravenous administration apparatus be replaced at least once every 24 hours.

Physicochemical studies have shown that the container and solution can withstand freezing.


1. Identify Two Ports (See Figure A).



2. To Add Medication


Remove additive port closure: hold container below additive port and grasp cap between thumb and forefinger then flip cap upward (**See Figure B**). **Swab exposed additive port.** Using a syringe with 18 gauge or smaller needle, insert cannula through resealable additive port and add desired drug. Mix thoroughly.

Note: Partial fill bags have been designed to accept an overflow of up to 50 mL.




3. To Attach Administration Set

To aseptically remove the set port closure: hold container below the set port and grasp the foil tab between the thumb and forefinger then pull the tab in two steps as shown in **Figure C Steps 1 and 2**.



4. Push spike through the diaphragm of the port (See **Figure D**). Hang container using hole on the lower flap. Prime set in accordance with the **Directions for Use** provided with the set in use.



When the container is to be used as a diluent and delivery system for intermittent intravenous administration of compatible drug additives, consult prescribing information for **INDICATIONS AND USAGE** of drug additives to be administered in this manner.

Warning: Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

PAB® containers can be safely transported in a standard 6-inch carrier through a pneumatic tube system that is well maintained and running properly.

B. Braun Medical Inc.

Bethlehem, PA 18018-3524 USA

1-800-227-2862

Y36-002-925 LD-224-5

PRINCIPAL DISPLAY PANEL - 25 mL Container Label

LOT

EXP

**0.9% Sodium Chloride
Injection USP**

NDC 0264-1800-36

REF S8004-5410

25 mL Partial Fill

in 100 mL PAB® Container

Each mL contains: Sodium Chloride USP 9 mg; Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5-7.0) Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/25 mL): Sodium 3.8 Chloride 3.8

Sterile, nonpyrogenic. Single dose container. For intravenous use only.

Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).

Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP.

Rx only

PAB is a registered trademark of B. Braun Medical Inc.

B. Braun Medical Inc.

Bethlehem, PA 18018-3524 USA

1-800-227-2862

Y94-003-353

LD-227-2

LOT

EXP



**0.9% Sodium Chloride
Injection USP**

NDC 0264-1800-36

REF S8004-5410

25 mL Partial Fill
in 100 mL PAB® Container

Each mL contains: Sodium Chloride USP 9 mg; Water for Injection USP qs
pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5-7.0) Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/25 mL): Sodium 3.8 Chloride 3.8

Sterile, nonpyrogenic. Single dose container. For intravenous use only.
Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with
pharmacist. When introducing additives, use aseptic techniques.
Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).

Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP. Rx only

PAB is a registered trademark of B. Braun Medical Inc.

Y94-003-353
LD-227-2

B | BRAUN

B. Braun Medical Inc.
Bethlehem, PA 18018-3524 USA
1-800-227-2862



PRINCIPAL DISPLAY PANEL - 50 mL Container Label

LOT

EXP

**0.9% Sodium Chloride
Injection USP**

NDC 0264-1800-31

REF S8004-5384

50 mL Partial Fill
in 100 mL PAB® Container

Each mL contains: Sodium Chloride USP 9 mg; Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5-7.0) Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/50 mL): Sodium 7.7 Chloride 7.7

Sterile, nonpyrogenic. Single dose container. For intravenous use only.
Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with
pharmacist. When introducing additives, use aseptic techniques.
Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).
Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP.

Rx only

PAB is a registered trademark of B. Braun Medical Inc.

B. Braun Medical Inc.

Bethlehem, PA 18018-3524 USA

1-800-227-2862

Y94-003-339

LD-226-A-2

LOT

EXP



0.9% Sodium Chloride Injection USP

NDC 0264-1800-31
REF S8004-5384

50 mL Partial Fill
in 100 mL PAB® Container

Each mL contains: Sodium Chloride USP 9 mg; Water for Injection USP qs
pH adjusted with Hydrochloric Acid NF
pH: 5.5 (4.5-7.0) Calc. Osmolarity: 310 mOsmol/liter
Electrolytes (mEq/50 mL): Sodium 7.7 Chloride 7.7

Sterile, nonpyrogenic. Single dose container. For intravenous use only.
Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with
pharmacist. When introducing additives, use aseptic techniques.
Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).
Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP. Rx only
PAB is a registered trademark of B. Braun Medical Inc.

Y94-003-339
LD-226-A-2

25

B | BRAUN

B. Braun Medical Inc.
Bethlehem, PA 18018-3524 USA
1-800-227-2862



PRINCIPAL DISPLAY PANEL - 100 mL Container Label

LOT
EXP

0.9% Sodium Chloride Injection USP

NDC 0264-1800-32
REF S8004-5264

100 mL Partial Fill
in 150 mL

PAB[®] Container

Each mL contains:

Sodium Chloride USP 9 mg

Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5-7.0)

Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/100 mL): Sodium 15.4; Chloride 15.4

Sterile, nonpyrogenic. Single dose container.

For intravenous use only. Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).

Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP.

Rx only

PAB is a registered trademark of B. Braun Medical Inc.

B. Braun Medical Inc.

Bethlehem, PA 18018-3524 USA

1-800-227-2862

Y94-003-337

LD-225-A-2

LOT

EXP

BARCODE

0.9% Sodium Chloride Injection USP

NDC 0264-1800-32

REF S8004-5264

100 mL Partial Fill

in 150 mL

PAB® Container

Each mL contains:

Sodium Chloride USP 9 mg

Water for Injection USP qs

pH adjusted with Hydrochloric Acid NF

pH: 5.5 (4.5-7.0)

Calc. Osmolarity: 310 mOsmol/liter

Electrolytes (mEq/100 mL): Sodium 15.4; Chloride 15.4

Sterile, nonpyrogenic. Single dose container.

For intravenous use only. Use only if solution is clear and container and seals are intact.

WARNING: Some additives may be incompatible. Consult with pharmacist. When introducing additives, use aseptic techniques. Mix thoroughly. Do not store.

Recommended Storage: Room temperature (25°C).

Avoid excessive heat. See Package Insert.

Not made with natural rubber latex, PVC or DEHP. Rx only

PAB is a registered trademark of B. Braun Medical Inc.

Y94-003-337
LD-225-A-2

—

25

50

75

B | BRAUN

B. Braun Medical Inc.
Bethlehem, PA 18018-3524 USA
1-800-227-2862



SODIUM CHLORIDE

sodium chloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0264-1800
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0264-1800-36	116 in 1 CASE	02/08/1978	
1		25 mL in 1 CONTAINER; Type 0: Not a Combination Product		
2	NDC:0264-1800-31	84 in 1 CASE	02/08/1978	
2		50 mL in 1 CONTAINER; Type 0: Not a Combination Product		
3	NDC:0264-1800-32	64 in 1 CASE	02/08/1978	
3		100 mL in 1 CONTAINER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017464	02/08/1978	

Labeler - B. Braun Medical Inc. (002397347)

Revised: 2/2018

B. Braun Medical Inc.